



Colonoscopy SuTab Bowel Preparation

PLEASE READ AND FOLLOW ALL INSTRUCTIONS CAREFULLY

Colonoscopy Preparation:

Your physician will need to provide a prescription to purchase SuTab preparation from your pharmacy.

You will need to have a very clean colon for the Gastroenterologist to examine your colon. This process requires taking strong laxatives which will cause significant diarrhea. You must follow the preparation instructions carefully to assure you will be cleaned out and to prevent dehydration. Occasionally despite your best efforts, the clean out will be insufficient and the procedure will have to be postponed. Your Gastroenterologist will make recommendations appropriate for your digestive system.

On the entire day before the procedure only liquids, no solid foods are allowed. NO solid foods are allowed for breakfast, lunch, or dinner. You may have any liquids such as water, juices, tea, coffee, soda, non-red sports drinks, and non-red Jell-O. Until 5 pm, you are permitted milk, ice cream with no solid particles, milk shakes and yogurt with no solid particles. After 5 pm, no milk products are allowed.

At 5 PM, begin the first half of the preparation:

- A. Open one bottle of 12 tablets.
- B. Fill the provided container with 16 ounces of water (up to the fill line).
- C. Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
- D. Approximately 1 hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- E. If you experience symptoms of (bloating, nausea, or cramping) pause the rate of drinking the additional liquid until symptoms diminish.

**The time to onset of action of the first dose of laxatives is very variable. It may not start for hours.

On the day of the procedure Approximately 6 hours prior to your scheduled arrival time, complete the second half of the preparation:

- A. open the second bottle of 12 tablets.
- B. Fill the provided container with 16 ounces of water (up to the fill line)
- C. Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
- D. Approximately 1 hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- E. Approximately 30 minutes after finishing the second container of water, fill the provided container again with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes. I
- F. If you experience symptoms of (bloating, nausea, or cramping) pause the rate of drinking the additional liquid until symptoms diminish.

You must stop drinking 3 hours before the procedure. If you have anything by mouth, including gum, in the three hours before the procedure, the procedure may be rescheduled.

No illicit substances 24 hours prior to the procedure and no alcohol or marijuana products for 12 hours prior. All patients will be assessed for signs of impairment on the day of the procedure and if there is any concern, the procedure will be cancelled.

If you take heart or blood pressure medications, take your pills three hours before the procedure with water (unless advised differently by your doctor).

Note: Please leave all jewelry and valuables at home. Optum Medical is not responsible for the loss of any patient's property. Do not wear contact lenses.